

# Transport and Magnetic properties of $\text{PrCoIn}_5$

Abebe Kebede\* and Terrel Dial

*Department of Physics*

*North Carolina Agricultural and Technical State University and  
1601 East Market Street, Greensboro, NC 27411*

(Dated: May 28, 2009)

Structural, electrical and magnetic measurements of 115 single crystals of  $\text{PrIn}_5$  are reported. It has a tetragonal structure and has slightly lower cell volume than its isomorphous counterpart  $\text{CeCoIn}_5$ . The resistivity saturates for  $T \geq 10\text{K}$ . Analysis of the resistivity for  $10\text{K} < T < 60\text{K}$  indicates a regular fermi liquid behavior. It does not exhibit superconductivity down to  $T \sim 1\text{K}$ . The magnetic susceptibility analysis yielded the moment to be  $4.00\mu_B$  indicating that the magnetism of  $\text{PrCoIn}_5$  is dominated by  $\text{Pr}^{3+}$  free ions with some admixture of the magnetic moment of the Co sublattice. The paramagnetic Curie temperature  $\theta \sim 40\text{K}$ . At low temperatures the susceptibility follows a broad maximum around  $T_N \sim 14.5\text{K}$ , and increases as the temperature is lowered. The disappearance of superconductivity for  $T > 1\text{K}$  is attributed to chemical pressure effects and magnetic pair breaking.

PACS numbers: 74.70.Tx, 74.25.Bt, 74.62.Bf

## I. INTRODUCTION

A typical metal is expected to exhibit fermi liquid behavior in that at low temperatures the electronic specific heat coefficient,  $\gamma$  and the magnetic susceptibility,  $\chi$  are constants, and the resistivity,  $\rho = \rho_0 + AT^2$ . Instead the normal state properties of  $\text{CeCoIn}_5$  are characterized by a non fermi liquid behavior where  $C/T \sim -\ln T$  and  $\rho = \rho_0 + AT^n$  ( $n < 2$ ) and  $\chi \sim T^{-n}$  ( $n < 1$ ). Its superconducting transition temperature  $T_c \sim 2.3\text{K}$  is significantly influenced by application of hydrostatic pressure and chemical substitution. In the system  $\text{CeMIn}_5$  ( $M = \text{Co, Rh, Ir}$ ), it has been shown that the superconductivity can be induced by tuning Neel temperature,  $T_N$  to zero. Substitution of In with Sn[1] and Ce with La[2] produced pair breaking effects attributed to chemical pressure[3]. In this work we synthesized good quality  $\text{PrCoIn}_5$  single crystals to study its structure, thermal and magnetic properties. The structurally similar  $\text{PrCoIn}_5$  is non superconducting, its resistivity saturates below  $10\text{K}$ . These differences in properties may allow researchers to study the evolution from superconducting non fermi liquid ground state exhibited by  $\text{CeCoIn}_5$  to a non superconducting fermi liquid state exhibited by  $\text{PrCoIn}_5$ . Single crystals of  $\text{PrCoIn}_5$  were synthesized from indium flux by combining stoichiometric amounts of Pr and Co with excess indium. The charge is put inside an alumina crucible, which is sealed in an evacuated quartz cylinder. It is then heated to  $1150\text{C}$  at the rate of  $2\text{C}$  per minute, followed by cooling to  $750\text{C}$  at  $3\text{C}/\text{minute}$ , rapid cooling to  $450\text{C}$  at  $2\text{C}/\text{minute}$ . The charge is then removed, and quickly put in centrifuge to remove the excess flux. Each good crystal is etched

TABLE I: Lattice parameters of  $\text{PrCoIn}_5$

Sample	$a(\text{\AA})$	$b(\text{\AA})$	$c(\text{\AA})$	$c/a$	Cell volume ( $\text{\AA}^3$ )
$\text{PrCoIn}_5$	4.61	4.61	7.50	1.62	161.09
$\text{CeCoIn}_5$	4.62	4.62	7.56	1.63	161.36

with concentrated hydrochloric acid for several hours and then it is rinsed with ethanol in ultrasonic environment. The resulting crystals are mostly plates with dimensions  $2\text{mm} \times 2\text{mm} \times 1\text{mm}$ .




FIG. 1: x-ray powder diffraction patterns of  $\text{PrCoIn}_5$

The lattice parameters were obtained from single crystal x-ray diffraction pattern with  $\text{Mo-K}\alpha$  radiation. The x-ray intensity as a function of  $2\theta$  is shown in Figure 1 above. The reflections were indexed for the tetragonal unit cell. Shown on Table 1 are the lattice parameters and the cell volumes of  $\text{PrCoIn}_5$  and  $\text{CeCoIn}_5$ . The  $c/a \sim 1.62$  clearly indicates that the structure of  $\text{PrCoIn}_5$  is tetragonal and it is structurally isomorphous with  $\text{CeCoIn}_5$ .

## II. RESISTIVITY OF $\text{PRCOIN}_5$

Resistivity was measured using a standard 4-wire technique at temperatures ranging from  $300\text{K}$  to  $1.8\text{K}$  in helium cooled cryostat. Excitation currents of

\*Electronic address: abkebede@gmail.com;  
URL: <http://sirius-c.ncat.edu/>

500 $\mu A$  – 1000 $\mu A$  were applied parallel to the c-axis at frequencies of 16 Hz. Each sample was mounted on a resistivity puck with GE varnish. #40 gauge copper wires were attached to the samples with silver epoxy. The magnetic susceptibility and magnetization measurements were carried out with a PPMS system at NHMFL-Los Alamos.




FIG. 2: Resistivity versus Temperature of PrCoIn<sub>5</sub>

As shown in Figure 2 the resistivity shows metallic behavior above about 150K. Since for typical metals such as copper, the deviation from linearity begins at  $T \sim 0.7\theta_D$ . We estimated  $\theta_D \sim 241K$ . Below 150K the data follow a broad maximum near  $T \sim 72K$  accompanied with slow decrease as the temperature is lowered. The broad maximum is attributed to crystalline electric field effects. There is no evidence of superconductivity within the accessible temperature  $T > 1K$ . Shown in Figure 3 is our fit of the data for  $10K < T < 60K$  using a polynomial of order five. Assuming that the contribution of the higher order terms diminishes as the temperature is lowered, we retained the fit parameters,  $\alpha = 0.034\mu\Omega-cm$ ,  $\beta = -0.005\mu\Omega-cm/K$  and  $A = 4.94 \times 10^{-4}\mu\Omega-cm/K^2$ . While the negative linear term can be an artifact of the fit, the  $T^2$  term is suggestive of a normal fermi liquid behavior. Extrapolated from the high temperature linear component we obtain the residual resistivity  $\rho_0 \sim 0.021\mu\Omega-cm$  and the residual resistivity ratio ( $RRR = \rho_{RT}/\rho_0$ ) was 200. This ratio indicates that this crystal is of reasonable quality. As shown in Figure 4 the resistivity saturates for  $T < 10K$  with  $\rho_{sat} \sim 0.034\mu\Omega-cm$ .  $\rho_{sat}$  is lower than the residual resistivity,  $\rho_0$ . Whether or not the origin of this saturation is of fundamental nature or caused by inclusions and surface effects is to be determined.




FIG. 3: Low Temperature Resistivity of PrCoIn<sub>5</sub>




FIG. 4: Saturation of the Resistivity versus Temperature of PrCoIn<sub>5</sub>

### III. MAGNETIC SUSCEPTIBILITY OF PRCOIN<sub>5</sub>

Shown in Figure 5 is the magnetic susceptibility of PrCoIn<sub>5</sub> measured at 1kOe. The data showed a rapid upturn at low temperature. The inset shows the susceptibility below 30K, the broad maximum near  $T_N \sim 14.5K$ . This structure coincides with the resistivity saturation for  $T < 10K$  and we believe it is associated with antiferromagnetic ordering of the Pr ions. While similar behaviour has observed in other Pr based alloys such as PrBa<sub>2</sub>Cu<sub>3</sub>O<sub>7</sub>, a non superconducting cuprate[4] and Pr<sub>3</sub>In[5] the presence of multiphases cannot be ruled out. Shown in Figure 6 is the plot of the inverse susceptibility,

TABLE II: Physical parameters of PrCoIn<sub>5</sub>

PrCoIn <sub>5</sub>	$T_K(K)$	$T_c(K)$	$T_N$	$\theta(K)$	$\mu(\mu_B)$
	72	0	14.5K	-40	4.00

$\chi^{-1}$  versus temperature.




FIG. 5: Magnetic Susceptibility versus Temperature of  $\text{PrCoIn}_5$  at  $H=1\text{kOe}$




FIG. 6: Inverse Magnetic Susceptibility versus Temperature of  $\text{PrCoIn}_5$  at  $H=1\text{kOe}$

At high temperatures ( $T \geq 100\text{K}$ ),  $\chi^{-1}$  is linear and follows Curie-Weiss behavior. We obtained effective moment of  $4\mu_B$ . This value is larger than the  $3.73\mu_B$  that is expected for a free  $\text{Pr}^{3+}$ . The higher magnetic moment could be attributed to an admixture of moments from both Co ions and  $\text{Pr}^{3+}$ . The Co sub lattice can be magnetized because of f-d exchange as in  $\text{RCO}_3$ [6]. Obviously the moment is dominated by  $\text{Pr}^{3+}$  in the  $J = 4$  Hunds rule ground state indicating that the  $4f$  electrons are almost localized within the  $\text{Pr}$  atoms. Similar results are obtained by others except that our analysis shows a lower paramagnetic Curie temperature  $\theta \sim -40\text{K}$  compared to  $-56\text{K}$ [7].

In summary we prepared clean single crystals of  $\text{PrCoIn}_5$ . The structure is tetragonal it has a  $c/a \sim 1.62$  slightly smaller volume than its counterpart  $\text{CeCoIn}_5$ . Superconductivity is not observed down to  $1\text{K}$ . In addition the behavior of the low temperature resistivity appears to have originated from a fermi liquid behavior. Pair breaking phenomenon were observed when  $\text{CeCoIn}_5$  is alloyed by the non magnetic La further demonstrating unconventional superconductivity in this system[2].  $\text{PrCo}_5$  has slightly lower ( $C/a \sim 1.62$ ) than that of  $\text{CeCoIn}_5$  ( $C/a \sim 1.63$ ). Whether or not this difference has such detrimental effect on superconductivity is not determined. However the high value of the magnetic moment and the presumed magnetic order below  $T_N \sim 14.5\text{K}$  may have led to magnetic pair breaking.

## Acknowledgments

This work was supported by the NASA MUCERPI2003 under the Grant NNG04GD63G. We are grateful to our collaborators National High Magnetic Field Laboratory-Los Alamos.

## IV. REFERENCES

- [1] E. D. Bauer, C. Capan, F. Ronning, R. Movshovich, J. D. Thompson, and J. L. Sarrao Physical Review Letters 94, 047001 (2005)
- [2] C. Petrovic, S. L. Budko, V. G. Kogan, and P. C. Canfield Physical Review B 66, 054534 (2002)
- [3] G. Sparna, R. Borth, E. Lengyel, P.G. Pagliuso, J.L. Sarrao, F. Steglich and J.D. Thompson Physica B 312313 (2002) 138139
- [4] A. Kebede, C. S. Jee, J. Schwegler, J. E. Crow, T. Mihalisin, G. H. Myer, R. E. Salomon, and P. Schlottmann M. V. Kuric, S. H. Bloom, and R. P. Guertin Phys. Rev. B 40, 4453-4462 (1989)
- [5] A. D. Christianson, J. M. Lawrence, J. L. Zarestky, H. S. Suzuki, J. D. Thompson, M. F. Hundley, J. L. Sarrao D. Antonio and A. L. Cornelius PHYSICAL REVIEW B 72, 024402 (2005)
- [6] I.Yu. Gaidukova, S.A. Granovsky, A.S. Markosyan, V.E. Rodimin Journal of Magnetism and Magnetic Materials 301 (2006) 124130
- [7] Nguyen Van Hieu1, Hiroaki Shishido, Arumugam Thamizhavel1, Rikio Settai1, Shingo Araki1, Yasuo Nozue1, Tatsuma D. Matsuda, Yoshinori Haga, Tetsuya

Takeuchi, Hisatomo Harima<sup>4</sup> and Yoshichika Onuki,J.  
Phys. Soc. Jpn. 74 (2005) pp. 3320-3328